

The Problem: What System to Plant?

- There is great disparity of opinion on the optimum planting density.
- Some growers plant 200-300 trees/acre on semidwarfing rootstocks with Central Leader.
- Most growers plant 500-1000 trees/acre on dwarfing rootstocks with some version of Vertical Axis.
- A few growers plant 2000 trees/acre on dwarfing rootstocks with Super Spindle.

Viable Orchard Systems in NY

• Tree		ensity	Spacing			
•System	(trees/acre)	(trees/ha)	(ft.)	Rootstocks		
•Slender Pyramid	340	840	8' x 16'	M.26, G.30,G.935		
•Vertical Axis	622	1538	5' x 14'	M.9, G.41,G.11		
•Slender Axis	908	2244	4' x 12'	M.9, G.41,G.11		
•Tall Spindle	1320	3262	3' x 11'	M.9, G.41,G.11		
•Super Spindle	2178	5382	2' x 10'	M.9, G.41,G.11		


Slender Spindle/M.9




Triple Row Slender Spindle/M.9




Geneva Y-trellis/M.26






USA-Vertical Axis/G.202

V-Slender Spindle/M.9




Gala Super Spindle/M.9 (Second Leaf)






Tall Spindle/M.9
(Note high branches that do not need to be tied up.

Braeburn Tall Spindle/M.9 (Third Leaf)



Tall Spindle/M.9





The Geneva Planting Systems Trial


(Planted 1997)

System	Spacing
Slender Pyramid/M.7	10' X 18'
Slender Pyramid/M.26	8' X 16'
Vertical Axis/M.9	7' X 15'
Vertical Axis/M.9	6' X 14'
Slender Axis/M.9	5' X 13'
Slender Axis/M.9	4' X 12'
Tall Spindle/M.9	3' X 11'
Super Spindle	2' X 10'










Conclusions

- Tree density had a strong influence on tree size, yield per hectare and light interception.
- The very high tree densities achieved a high yield of 50 Mt/ha by the fourth year and had a cumulative yield 3X the lowest densities.
- Tree shape had no effect on tree size or yield.

		Tree	Yield/Acre (bu.)			
Variety	System	Den.	2nd	3rd	4th	Cum
•McIntosh	Central Leader	218	0	1	24	25
•	Slender Pyramid	444	0	37	113	150
•	Vertical Axis	726	2	239	327	566
•	Tall Spindle	1307	13	351	771	1135



With traditional fruit prices, profitability over 20 years increases with increasing tree density up to a density of 1,000-1,100 trees/acre.

Can you afford to replant and what density should you choose?

- If you receive a farm gate fruit price of at least \$5.50/bu., have land or can buy land for up to \$5,000/acre, can purchase trees for a price of \$5.32, and have access to capital at 5% then you can afford to replant.
- The greatest profitability will be with the Slender Axis system (4' in-row spacing=908 trees/acre) or the Tall Spindle system (3' in-row spacing=1320 trees/acre)
- A 4 wire trellis and feathered trees are essential components of the system.

The Tall Spindle and Slender Axis Systems

- 3-4' in-row spacing
- •10-12' between rows
- •10' tall
- no permanent branches
- highly feathered trees
- minimal pruning at planting
- feathers tied below horizontal at planting.





Tall Spindle

Rootstock Selection

- M.9 T337 on vigorous soils or with vigorous scions, Nic29 or Pajam 2 on replant soil or with weak scions)
- B.9 or O.3 are alternatives.
- G.41 or G.11 may be better.

Recommended Tree Type for the Tall

Tree Spacing of the Tall Spindle

- In-row spacing of 3' for weak growing varieties (Honeycrisp, Delicious, Empire, Macoun, Gala).
- In-row spacing of 4' for vigorous varieties and tip bearing varieties. (McIntosh, Fuji, Cortland, Jonagold, Gingergold).
- Between row spacing should be 12-13' on slopes and 10-11' on level ground.

Pruning after Planting of Tall Spindle

Large caliper highly feathered trees (8-15 feathers).

- Do not head the leader.
- Remove all feathers below 24".
- Remove feathers that are larger than 2/3 diameter of leader.
- Do not tip the feathers.

Medium Caliper poorly feathered trees or whips.

- Head the leader at 60".
- Remove feathers larger than 2/3 diameter of leader.
- If there are less than 3 good feathers remove them using a bevel cut.
- Score above every other bud along leader from 24"-42" high.








Tree Training During the First Summer

- Pinch side shoots in upper 1/4 of leader when shoots are 4-5" long.
- Re-pinch side shoots in upper 1/4 of leader when regrowth is 4-5" long.
- Tie leader to support system.
- Tie down 4-5 lower branches below horizontal at planting or in July to induce flowering.



Pruning Year 2

- Do not head leader.
- Do not head feathers
- Remove side branches that compete with leader using a bevel cut.
 - Remove narrow angled branches.
 - Remove scaffolds that are larger than 2/3 diameter of leader.
 - Remove side branches that are longer than 2'.
- Remember "large branches create large trees"





Early Cropping


- "The best way of restricting vegetative growth is to produce apples." (Kurt Werth)
- Cropping must begin:
 - In the second year with the Tall Spindle system.
- Cropping targets for the Tall Spindle
 - Year 11-5 fruits
 - Year 220 fruits
 - Year 3 40 fruits
 - Year 470 fruits
 - Year 590 fruits

Pruning Year 3-5

- Allow crop to bend the top.
- Limit height of tree only after top has bent by cutting leader to a weak fruitful side branch.
- Remove branches larger than 3/4 inch diameter.
- Remember "large branches create large trees"
- Shorten older, pendant branches to a weak side branch or spur.
- With Gala begin stubbing back pruning.



Pruning Yee.j 1wh0







Conclusions

- 1) The tall spindle or Slender Axis systems appear to be the most profitable systems.
- 2) High tree density gives high early yield.
- 3) Highly feathered trees are the key to the systems.
- 4) Minimal pruning at planting (No heading the leader or tipping the feathers at planting)
- 5) Branch angle management. Bending feathers below horizontal at planting induces early cropping and limits branch size.
- 6) Branch caliper management. Ruthless removal of large branches keeps trees manageable. "Large branches create large trees"
- 7) "The best way of restricting vegetative growth is to produce apples."