2019-2021 Brassica Cover Crops in Vegetable Rotations Trial

Dr. Heather Darby, UVM Extension Agronomist
Rory Malone, Scott Lewins, Dr. Vic Izzo, and Ivy Krezinski
UVM Extension Crop

© April 2022, University of Vermont Extension

2019-2021 BRASSICA COVER CROPS IN VEGETABLE ROTATIONS TRIAL

Dr. Heather Darby, University of Vermont Extension
heather.darby[at]uvm.edu

Brassica crops are an important crop on diversified vegetable farms in Vermont and the Northeast. Brassicas are becoming more popular as a cover crop due to their unique benefits to soil health. They provide ground cover and cover crop biomass rapidly in the fall, and brassicas with long taproots further break up soil compaction. Brassicas grown on vegetable farms include rutabaga, cabbage, cauliflower, kale, broccoli, Brussels sprouts, collards, and mustard greens. Despite the benefits, not all vegetable producers are quick to add brassica cover crops to their brassica cash crop rotations due to concern that pests and disease are likely to transfer from one brassica crop to another. An increased use of high tunnels has also created a To assess the risk of pest and disease carryover between brassica crops, the University of Vermont Extension Northwest Crop and Soils (NWCS) Team conducted a field trial at two locations over two full field seasons with fall-planted brassica cover crops, followed by a broccoli cash crop.

MATERIALS AND METHODS

The trial was conducted at

2020 Broccoli scouting period	12-Jun to 24-Jul	10-Jun to 25-Jul
2020 Cover crop scouting period	9-Sep to 7-Oct	3-Sep
2021 Broccoli scouting period	8-Jun to 19-Jul	17-

main effect was followed by pairwise comparisons and a significant interact was followed by examination of simple effects. Mean comparisons

significantly higher in the HGM brassica plots than the

Table 12. 2020-2021 Horticulture Research Farm Treatment Results, South Burlington, VT.

FB= flea beetle, ICW= imported cabbage worm, DBM= diamondback moth. Statistical significance is based on p<0.05.

Table 13. 2020-2021 Borderview Research Farm Treatment Results, Alburgh, VT.

FB= flea beetle, ICW= imported cabbage worm, DBM= diamondback moth. Statistical significance is based on p<0.05

Table 14. 2020-2021 Borderview Research Farm Significant Results by Treatment, Alburgh, VT.

Treatment	Fall 2020 Downy Mildew Means	Fall 2020 Flea Beetle Means
HGM	**0.0073[a]**	**0.0333[a]**

weather was warm and dry in both years of the study. Additionally, the fall cover crops in 2020 at the South Burlington site were only scouted on one date. This trial adds to the little research available on brassica disease carryover in the Northeast, and further research is needed.

For more information on pathogens, pests, and Integrated Pest Management, see the New England Vegetable Management Guide, available at: https://nevegetable.org/

References

New England Vegetable Management G