2020 Industrial Hemp Fiber Variety Trial



Dr. Heather Darby, UVM Extension Agronomist
John Bruce, Ivy Luke, and Rory Malone
UVM Extension Crops and Soils Technicians
(802) 524-6501

Visit us on the web: http://www.uvm.edu/nwcrops

© December 2020, University of Vermont Extension

Table 2. Hemp varieties evaluated in the industrial hemp fiber trial 2020, Alburgh, VT.

Variety	Days to maturity	Seed supplier
Bialobrzeskie	130-145	Bija Hemp
Henola	115-120	Bija Hemp
Hlukhivs'ki 51	120-125	Fiacre Enterprises
Hlesia	115-120	Fiacre Enterprises
Hliana	115-120	Fiacre Enterprises
Futura	129-134	UniSeeds/Seedway
Altair	100	UniSeeds/Seedway
Anka	110	UniSeeds/Seedway
Ferimon	140-145	UniSeeds/Seedway

On 23-Jun, the trial was fertilized with 200 lbs ac^{-1} of urea (46-0-0). Fertility amendments were based on soil test results indicating that no additional fertility was required for potassium or phosphorus. Select Max® was sprayed on the trial on 15-Jul at a rate of 16 oz ac^{-1} to limit weed pressure from grasses.

On 24-Jun, plant populations were recorded by counting the number of plants in a foot-long section of the row, two times per plot. Prior to mowing, three randomly selected plant heights were recorded on 7-Aug within each plot. Additionally, on the same day, wet weight harvest yields were calculated by sampling the hemp biomass within a 0.25m^2 quadrat. Harvest moisture was calculated by taking a subsample of hemp yield and drying it at 105 F until it reached a stable weight. On 10-Aug, five plants were selected at random from each plot and run through a custom-built decorticator (Image 1). While the stalks were still fresh, they were decorticated to separate the bast and hurd fibers. As the stalks passed between the two moving gears, hurd fiber broke away and dropped to a bucket placed underneath.



Image 1. Custom built decorticator, Alburgh, VT, 2017.

Table 4. The impact of variety on plant characteristics and harvest yield of industrial hemp fiber, Alburgh, VT, 2020.

Variety	Plant height	Harvest moisture	Dry plant weight	Dry matter yield	Harvest population	Harvest population	Bast fiber
	cm	%	lbs plant^{-1}				

Figure 1.

break down plant cell tissues and pectin that bind the fibers together. The speed of this process in most common cases is influenced by moisture and temperature directly impacting microbial activity which is responsible for breaking down plant material. Warm and moist conditions will encourage increased microbial activity and thus speed the retting process. Moisture is required for the process to occur but, depending on temperature,