

Lichen Monitoring

A relative epiphytic simplicity

Lichens & symbiotic respond t generally have diff effects of indicators Recent w communi northeast. commonl

At the Ve and abunc survey ear on Mansfi only folio: of sulfur s has varied *Allocetrar* species, *P.*

Data colle for 2002. communit

ely new inc ni lichen con n y of lichens n ure unique n ally. They e o environn er be descrip w erent liche s pollution a o : of regional e ork by the fa ties as indicat Since lichen y used in inter

rimont Mont k lance in ass oc chi of the L p field. Using n sel and fruti co ensitive liche from 11 to 1 *ia odkestanica*, *uncyella apyca*

ction in 2001 In addition, to ies.

oring eff unity con takes the

the plant ly exclus ital condi e of the e species tha und a nu air quality tional Fo ors of bo species c tifying p

ring Coc iation wi lots on M ethods de se lichen ns have t 8 species *Evernia lachensis*

will com these sit

or, cond n position multiple f world's tive y n tions are ny, recom in a forest mber of p in, th, u j res. Heal th, ege on differ t as ote, nija a

pe, at we th pac o for n Ma ve opp ed s, we hav ycep foun per lqat m, so, for was you

plate plo es, 22 NE

dy Wilmot and ment of Forests, F

acted by the Dep t and abundance or use as indicato

being a combina he atmosphere a h as light, moist ental conditions t edge). Lichens joint sources and per Ohio Valley th Monitoring; pr al air quality, gra ensitivity to sulfi ir pollution effec

sites, we are curr f the forest healt nsfield and 5 at I by the National F e thus far identif d, and have ocbu ion. The most co *pha* and *Hypogytona*

ts on Mount Mar

or, cond n position multiple f world's tive y n tions are ny, recom in a forest mber of p in, th, u j res. Heal th, ege on differ t as ote, nija a

pe, at we th pac o for n Ma ve opp ed s, we hav ycep foun per lqat m, so, for was you

plate plo es, 22 NE

Simonon s & Recre

ment of Fo measure f forest re

of a fung source of r and air qu given loca e been suc ustrial citi onnecticut am has be ts, and re; oxide and n forests.

y compili onitoring Brook, th st Health 46 species l across el only foun *a physode*

ld. The L ation

ar n s a rs' c tion s a re, at a hav in d Co ogr fier and s' c

ent h m ye fore led. rec um m.

nsfic ide

are invol

res st. Parks & Recrea io of for st health. The ale spous e to environme tal us, and an algae, grow ng nitation and moisture A ality. The lichen con ne tion (e.g. a dance for st pece s fully used to ma g es. They have also been ard in several Euro pa an are successful in using lic gional climatic gradients l surface, this characte is

ng paneline data on li he plots nontored. Whi e t us far 7 plots have be n Mcput print program, vt s from Mt. Mansfield Fi ever, in gradients. Sp ci d species thus far have t to. One regionally un on

ye Brook lichen surv y i ved in data collection or

at n uses relative n condi ns.

ving e. as such ey ur unity ca e stand v uld p round- el ec used al n coun s. g chen in the er lic is

ic n spec ille the goa to e comple f which sur ys ve spec s es rich s ve been: common

ey s plant f lichen

of for st health. The ale spous e to environme tal us, and an algae, grow ng nitation and moisture A ality. The lichen con ne tion (e.g. a dance for st pece s fully used to ma g es. They have also been ard in several Euro pa an are successful in using lic gional climatic gradients l surface, this characte is

ng paneline data on li he plots nontored. Whi e t us far 7 plots have be n Mcput print program, vt s from Mt. Mansfield Fi ever, in gradients. Sp ci d species thus far have t to. One regionally un on

ye Brook lichen surv y i ved in data collection or

Figure 1-3. Overstory tree health in the Lye Brook Wilderness Area. Tree health (Figure 2), and foliage transparency shows significant differences

at 2 elevations (baseline) for density (Figure 1) and dieback (figure 2) at 2 elevations (baseline) for density (Figure 1) and dieback (figure 2) at 2 elevations, “\*\*”

