Ready 2 Yield) soybean was obtained from Seedway, LLC (Hall, NY) for the trial. Soybeans were sprayed with

In this example, treatment C is significantly different from treatment A but not from treatment B. The difference between C and B is equal to 1.5, which is less than the LSD value of 2.0. This means that these treatments did not differ in yield. The difference between C and A is equal to 3.0, which is greater than the LSD value of 2.0. This means that the yields of these treatments were significantly different from one another.

RESULTS

Weather data was recorded with a Davis Instrument Vantage Pro2 weather station, equipped with a WeatherLink data logger at Borderview Research Farm in Alburgh, VT (T3). Overall, the season began cooler and wetter than normal but became hot and dry in the middle of the summer. The month of July brought above normal temperatures and little rainfall. The longest period without rainfall in July lasted 12 days. This dry period, which occurred around the time of pod formation, may have negatively impacted soybean plant growth and productivity. However, these environmental conditions were welcomed and provided the crop with crucial Growing Degree Days (GDDs) needed to reach maturity. The season had a total of 2400 GDDs accumulated May-

Table 4. Cover crop and soybean harvest characteristics, 2018-2019.

Cover crop treatment	Overwinters	Fall 2018 Spring 2019	Soybean harvest 2019		
		Dry matter yield	Yield at 13% moisture		Test weight



Figure 2. Fall biomass and soybean yield by cover crop mixture treatment, 2019.
Treatments that share a letter performed statistically similarly to one another. Soybean yields did not differ statistically across treatments