2023 GSR FERTILIZER TOMATO FIELD TRIAL

Dr. Heather Darby, University of Vermont Extension
heather.darby[at]uvm.edu

There is current interest in new fertilizer sources that optimize the use of natural resources while providing similar efficacy in crop fertilization and nutrient cycling as traditional fertilizers. GSR Solutions LLC (General Systems Research) produced an algae-based, naturally derived fertilizer. In this trial, the soil and leaf nitrate level and yields of tomato were evaluated to compare the GSR fertilizer to common nitrogen fertilizers during the 2023 field season.

MATERIALS AND METHODS

Tomato (var 'Galahad') seedlings were transplanted on 25-May into 10' x 5' plots. Plant spacing was 2' with 5 plants per plot (Table 1). The experimental design was a randomized complete block with four replicates. The soil type was Benson rocky silt loam with 3-5% slopes. Treatments included GSR's G3 fertilizer (3-0-0), Pro-Booster (10-0-0) by North Country Organics (Bradford, VT), Urea (46-0-0), hemp seed meal (6-1-1) produced at Borderview Farm (Alburgh, VT), and an untreated control. Soil amendments were applied at a rate to supply a total of 120 lbs N ac^{-1} (Table 2). Amendments were broadcast by hand and worked into the soil prior to transplanting.

Table 1. GSR tomato field trial specifics for Alburgh, VT, 2023.

from the center three plants of each plot on 30-Jun and 19-Jul. Thirty leaf petioles per plot were collected (10 petioles per plant), from the most recent mature leaves, into a composite sample and dried down prior to shipping. Samples were submitted to Dairy One Laboratory (Ithaca, NY) and analyzed for total nitrogen (%).

At harvest, the ripe tomatoes were picked from the center three plants in each plot, using the outside plants as buffers between treatments. Plots were harvested on six different occasions including 3-Aug, 9-Aug, 14-Aug, 21-Aug, 28-Aug, and 6-Sep. The number of ripe fruit and weight (lbs) were recorded at harvest.

Stand characteristics were analyzed using mixed model analysis using the mixed procedure of SAS (SAS Institute, 1999). Replications within the trial were treated as random effects, and treatments were treated as fixed. Treatment mean comparisons were made using the Least Significant Difference (LSD) procedure when the F-test was considered significant ($p<0.10$).

Table 3. Seasonal weather data for Alburgh, VT, 2023.

	May	Jun	Jul	Aug	Sep
Average temperature (°F)	57.1	65.7	72.2	67.0	63.7
Departure from normal	-1.28	-1.76	-0.24	-3.73	1.03
Precipitation (inches)	1.98	4.40	10.8	6.27	2.40
Departure from normal	-1.78	0.14	6.69	2.73	-1.27
Growing Degree Days (50-86°F)	303	483	712	540	449
Departure from normal	1.00	-41.0	17.0	-101	62.0

Based on weather data from a Davis Instruments Vantage Pro2 with WeatherLink data logger. Historical averages are for 30 years of NOAA data (1981-2020) for Burlington, VT.

DISCUSSION

This trial suggests slight differences between the GSR fertilizer and the other nitrogen fertilizers evaluated, as well as the control. Although, some data shows greater yields for the GSR on the 9-Aug harvest, it also shows fewer ripe tomatoes and weight on the 21-Aug harvest compared to the urea treatment. When comparing total pounds of tomatoes harvested, the GSR fertilizer did have a higher weight of fruit collected than both the Pro-Booster and hemp meal fertilizers, suggesting the GSR fertilizer may aid in more fruit development.

The soil nitrate levels for the fertilizer treatments were not statistically different, suggesting that they all had a similar nitrogen uptake. However, when looking at the leaf nitrate levels, GSR had the lowest levels when the samples were taken on 30-Jun, suggesting that more time is needed for nitrogen to become available to the plant. By the time of the second leaf nitrate sampling on 19-Jul, concentrations were similar which suggests nitrogen levels were similarly available among the different treatments. It is important to remember that these results only