2011 VERMONT BROWN MID-RIB CORN VARIETY TRIAL

Brown mid-rib (BMR) corn has a lower lignin content than other silage corn varieties, and many growers are interested in this type of corn for its higher digestibility and potential to yield more milk than conventional silage corn. With BMR's high fiber digestibility it may help increase dry matter intake of forage and may reduce purchase of concentrates. In 2011, the University of Vermont Extension conducted a research trial to evaluate the yield and quality of ten BMR varieties. While the information presented can begin to describe the yield and quality performance of BMR corn varieties in this region, it is important to note that the data represents results from only one season and one location. Compare other hybrid performance data before making varietal selections.

MATERIALS AND METHODS

The BMR corn silage variety trial was conducted at Borderview Farm in Alburgh, Vermont. The soil was a Benson rocky silt loam, and the previous crop was corn. The experimental design was a randomized block with three replications. The plot size was 5' x 50'. The seedbed was prepared with spring plowing and harrowing and finished with a spike-toothed harrow. The corn was planted on May 25 at a rate of 34,000 seeds per acre with a John Deere 1750 four-row corn planter. A 10-20-20 starter fertilizer was applied at 260 lbs per acre, and additional fertilizer (ammonium sulfate) was applied as a sidedress at 99 lbs of actual N per acre. On June 15, Lumax® (S-Metolachlor, Atrazine, and Mesotrione) was applied at a rate of 2 quarts per acre. Trial agronomic information is summarized in Table 1.

Table 1. Agronomic information for the 2011 BMR Variety Trial at Borderview Farm.

Location	Borderview Farm – Alburgh, VT
Soil type	Rocky Benson silt loam
Previous crop	Corn
Tillage operations	Spring plow, harrow, spike-toothed harrow
Plot size (ft.)	5' x 50'
Replicates	3
Seeding rate	34,000 seeds ac^{-1}
Row width (in.)	30
Planting date	25-May
Starter fertilizer	260 lbs ac^{-1} of 10-20-20
Topdress fertilizer	99 lbs N ac^{-1} at V6 stage
Herbicide	15-Jun, Lumax®, 2 qt ac^{-1}
Harvest dates	26-Sep and 7-Oct

The seed for this trial was donated by two participating seed companies, Mycogen and Seedway, LLC, whose contact information is listed in Table 2. Varieties ranged from 90-109 days in relative maturity.

Table 4. Impact of variety on selected yield and quality characteristics of BMR corn, Alburgh, VT, 2011.

Variety	Harvest date	Population	Yield at 35% DM	DM at harvest	Forage quality characteristics						Milk per	
					CP	ADF	NDF	dNDF	Starch	NEL	ton	acre
		plants/ac	tons/ac	%	%	%	%	%	%	Mcal/lb	lbs	lbs
5555BMR	9-Sep	29200	16.3	35.3	7.9	22.4	38.9*	71.6*				



Figure 2. ADF and NDF concentrations of ten BMR corn varieties. Treatments with the same letter did not differ significantly (p=0.10).

Figure 3. Digestible NDF and milk per ton by variety. Treatments with the same letter did not differ significantly (p=0.10; compare lower-case letters for dNDF and capital letters for milk per ton).

Milk per ton, an indicator of silage quality, was highest in the varieties F2F383 and 5555BMR. This was not significantly higher than RST100, F2F387, or F2F488. There was no significant difference in milk per acre, which indicates both silage quality and yield.