2012
Vermont Organic Corn Silage

©December 2012, University of Vermont Extension

levels greater than 30% are not considered to affect energy content, and might in fact have a negative impact on digestion. Starch levels vary from field to field, depending on growing conditions and variety.

The silage performance indices of milk per acre and milk per ton were calculated using a model derived IURPWKHVSUHDGVKHHWHQWLWOHG0,/.GHYHORSHGEUHVHDUFKHUVDWWKH8QLYHUVLWRI:LVF
Milk per ton measures the pounds of milk that could be produced from a ton of silage. This value is generated by approximating a balanced ration from corn silage that meets animal energy, protein, and fiber needs. The value is based on a standard cow weight and level of milk production. Milk per acre is calculated by multiplying milk per ton by silage dry matter yield. Therefore, milk per ton is an overall indicator of forage quality and milk per acre an indicator of forage yield and quality. Milk per ton and milk per acre calculations provide relative rankings of forage samples, but should not be considered as predictive of actual milk responses in specific situations for the following reasons:

1) Equations and calculations are simplified to reduce inputs for ease of use,
2) Farm to farm differences exists,
3) Genetic, dietary, and environmental differences affecting feed utilization are not considered.

temperature scale, was 2,717 days in Alburgh and 2,260 days in Randolph. For the growing season, the accumulated GDD was above average by 324 GDD in Alburgh and 326 GDD in Randolph.

Table 4. 2012 Weather data for Alburgh and Randolph, Vermont.

Alburgh, VT	May	June	July	August	September	October
Average temperature (°F)	60.5	67.0	71.4	71.1	60.8	52.4
Departure from normal	4.10	1.20	0.80	2.30	0.20	4.20



Figure 1. Yield and protein of organic corn silage varieties, Alburgh, VT, 2012. Varieties with the same letter above the bar are not significantly different from each other.

Figure 2. Milk performance of corn silage varieties, Alburgh, VT, 2012. Dotted lines indicate milk per ton and milk per acre averages.

Randolph Yield and Quality

Corn silage grown in Randolph, VT had an average population of 31,000 plants acre^{-1} and an average yield of 16.4 tons acre^{-1} (Table 6). There were no significant differences in population, yield, or quality characteristics for the corn grown in Randolph (Table 6; Figure 3). The corn was harvested outside of the optimum moisture range of 60 to 70%. The average moisture content of the trial was 76% moisture (data not shown), which may account for some of the lack of differences noted between varieties.

Table 6. Organic corn silage yield and quality, Randolph, Vermont, 2012.

	Pop	Yield	Forage Quality Characteristics									Milk per	
	plants	tons	CP	ADF	NDF	dNDF	Starch	NEL	TDN	NFC	NSC	ton	acre
Variety	ac^{-1}	ac^{-1}	%	%	%	%	%	Mcal lb^{-1}	%	%	%	lbs.	lbs.
C912	30119	22.1	8.0	28.4	47.3	51.4	29.3	0.740	70.8	41.4	31.0	2735	21207
Vp2p78	32110	18.1	7.3	30.7	49.6	52.9	28.2	0.725	69.9	39.5	29.4	2737	17267
VO44-86	35346	17.0	6.7	27.8	44.9	55.1	31.7	0.750	72.1	44.8	33.7	2898	17175
23A71	29621	17.0	7.2	28.5	46.9	53.2	31.0	0.740	71.3	42.6	32.2	2824	16829
B915	31861	16.9	7.7	29.5	48.6	52.9	29.0	0.735	70.6	40.4	30.1	2790	16560
C714	31612	16.8	7.2	29.5	48.2	53.9	29.1	0.740	71.0	41.5	30.7	2829	24842
25A16	30865	16.1	6.9	27.8	45.7	54.8	32.1	0.755	72.4	44.3	33.7	2895	16250
Vp3P26	32359	15.9	7.4	28.2	47.1	54.3	29.8	0.750	72.0	42.1	31.6	2879	16049
3G33	27878												

The figures below display the relationship between milk per ton and milk per acre. The dotted lines dividing the figures into four quadrants represent the mean milk per ton and milk per acre for the location. Therefore, varieties that fall above and to the right of the lines performed above average and hybrids below or to the left of the lines performed below average. It is important to remember that varieties at the Randolph site did not differ significantly in yield or quality. Milk per ton measures the pounds of milk that could theoretically be produced from one ton of silage. Milk per acre is calculated by multiplying the milk per ton value by silage dry matter yield. Thus, milk per ton is an overall indicator of forage quality and milk per acre is an indicator of forage yield and quality. C714 (American Organic) was the only YDULHWWKDWIHOOLQWKHTXDGUDQWDERYHDYHUDJHDWERWKORFDWLRQV (Figure 2 and 4).

Figure 4. Milk performance of corn silage varieties, Randolph, VT, 2012. Dotted lines indicate milk per ton and milk per acre averages