THE EFFICACY OF SPRAYING FUNGICIDES TO CONTROL FUSARIUM HEAD BLIGHT INFECTION IN SPRING MALTING BARLEY

Dr. Heather Darby, University of Vermont Extension
Heather.Darby[at]uvm.edu

Public interest in sourcing local foods has extended into beverages, and the current demand for local brewing and distilling ingredients is quickly increasing. One new market that has generated interest of both farmers and end-users is malted barley. This only stands to reason since the Northeast alone is home to over 180 microbreweries and 37 craft distillers. Until recently, local malt was not readily available to brewers or distillers. However, a rapid expansion of the fledgling malting industry will hopefully give farmers new markets and end-users hope of readily available malt. To date, the operating maltsters struggle to source enough local grain to match demand for their product. In addition to short supplies, the local barley that is available often does not meet the rigid quality standards for malting. One major obstacle for growers is *Fusarium* head blight (FHB) infection of grain. This disease is currently the most important disease facing organic and conventional grain growers in the Northeast, resulting in loss of yield, shriveled grain, and most importantly, mycotoxin contamination. A vomitoxin called Deoxynivalenol (DON) is considered the primary mycotoxin associated with FHB. The spores are usually transported by air currents and can infect plants at spike emergence through grain fill. Eating contaminated grain greater than 1ppm poses a health risk to both humans and livestock.

Fungicide applications have proven to be relatively effective at controlling FHB in other barley growing regions. Limited work has been done in this region on the optimum timing for a fungicide application to barley specifically to minimize DON. In addition, there are limited studies evaluating organic approved biofungicides, biochemicals, or biostimulants for management of this disease. In April 2017, the UVM Extension Northwest Crops and Soils Program initiated year four of a spring barley fungicide trial to determine the efficacy and timing of fungicide application to reduce FHB infection on cultivars with varying degrees of disease susceptibility.

MATERIALS AND METHODS

A field experiment was established at the Borderview Research Farm located in Alburgh, VT on 27-Apr to investigate the effects of cultivar resistance, fungicide efficacy, application timing on FHB and DON infection in spring malting barley. The experimental design was a randomized complete block, with a split-plot arrangement of cultivar as the whole-plot and fungicide+timing treatments as the sub-plots. The main plot of cultivar included Rasmussen, a 6-row malting barley that is a FHB susceptible variety, and Conlon, a 2-row malting barley with moderate FHB resistance.

Location	Borderview Research Farm Alburgh, VT
Soil type	Benson rocky silt loam

When the barley reached 50% spike emergence (22-Jun), plots were sprayed with the fungicide treatments (Table 2). The adjuvant 'Induce' was added to the Prosaro and Caramba applications at a rate of 0.125%. All but one plot (Control) of each cultivar was inoculated 24 hours (23-Jun), after the heading treatment was applied, with a spore suspension (40,000 spores/ml) consisting of a mixture of isolates of *Fusarium graminearum* endemic to the area. The *Fusarium graminearum* spores were multiplied and harvested using the 'Gz conidial suspension inoculum protocol'. Four days after the heading application (26-Jun) plots not previously treated with a fungicide were sprayed with the fungicides treatments except for the control and *Fusarium graminearum* only plots (Table 2). Water was applied at the same rate as the fungicides to the control plots and to those that were only inoculated with *Fusarium graminearum.* The application was made using a Bellspray Inc. Model T4 backpack sprayer. This model had a carbon dioxide pressurized tank and a four-nozzle boom attachment. It sprayed at a rate of 10 gallons per acre. Below is a list of the treatment materials evaluated in this trial. Descriptions have been provided from manufacturer information.

Actinovate® (EPA# 73314-1) is a biological fungicide (0.0371% Streptomyces lydicus WYEC 108) that suppresses and controls root rot, damping-off fungi and foliar fungal pathogens. Its active ingredient is a patented bacterium that grows around the root system (when soil drenched) and foliage of the plant (when sprayed on) while using several novel modes of antifungal action to protect plants.

Caramba® (EPA# 7969-246) fungicide is a highly effective fungicide containing the active ingredient metconazole, resulting in significant yield protection and reductions of deoxynivalenol (DON) levels in grain. It is not only effective on head scab, but provides control of late-season foliar diseases as well.

ChampION® (EPA# 55146-1) is a 77% copper hydroxide-based, broad-spectrum fungicide for disease control. When copper hydroxide is mixed with water, it releases copper ions, which disrupt the cellular proteins of the fungus. This product is approved for use in organic production systems.

Prosaro® (EPA# 264-862) fungicide provides broad-spectrum disease control, stops the penetration of the fungus into the plant and the spread of infection within the plant and inhibits the reproduction and further growth of the fungus.

SONATA® (EPA# 69592-13) fungicide provides excellent control of powdery mildews and rusts. Based on a patented strain of Bacillus pumilus (QST 2808), SONATA is an excellent fit for integrated disease

management programs. SONATA contains a unique, patented strain of Bacillus pumilus (QST 2808) that produces an antifungal amino sugar compound that inhibits cell metabolism. SONATA also creates a zone of inhibition on plant surfaces, preventing pathogens from establishing on the plant.

Table 2. Plot treatments-fungicide application dates and rates.

Treatments	Heading application	4 days after heading application	Application rate
	date	date	
Control	22-Jun	26-Jun	Water
Fusarium graminearum	23-Jun		

All fungicide+timing treatments



Figure 1. The impact of application timing and fungicide on barley yield and DON concentration.
Treatments with the same letter did not differ significantly.

Impact of Variety

There were significant differences in the average FHB plot severity, infected head severity, and incidence of FHB infection between malting barley varieties (Table 6). The variety Conlon had the lowest average FHB plot severity (5.23%), infected head severity (11.2%), and incidence of FHB infected heads (42.3%).

Table 6. The impact of malting barley variety of FHB incidence and severity.

Variety	Average FHB severity	Average FHB infected head severity	Incidence FHB of infected heads
	%	%	%
Conlon	5.23	**11.2**	**42.3**
Rasmussen		14.9	51.8
LSD (0.10)	1.57	1.69	5.82
Trial Mean	6.75	13.1	47.1

Values shown in **bold** are of the highest value or top performing.

The malting barley varieties were significantly different in harvest moisture, test weight, hundred-kernel weight, and DON (Table 7, Figure 2). Rasmussen had the lowest harvest moisture (14.2%). Both varieties had moistures above 14% and therefore had to be dried down for storage. Conlon had the highest test weight of 44.5 lbs bu^{-1}, heaviest hundred-kernel weight (5.00 g) and the lowest DON concentration (4.53 ppm). Neither of the varieties achieved industry standards for test weight of 48 lbs bu^{-1}, and both had DON concentrations above the FDA 1 ppm recommendation. Varieties did not differ in yield (Figure 2).

Table 7. The impact of malting barley variety of quality and yield.

Variety	Harvest moisture	Test weight	Hundred kernel weight	Yield @13.5% moisture	DON
	%	lbs bu^{-1}	grams	lbs ac^{-1}	ppm
Conlon	14.8	**44.5**	**5.00**	2828	**4.53**
Rasmussen	**14.2**	43.9	4.11	**2854**	8.29
LSD (0.10)	0.53	0.52	0.06	NS	0.54

DISCUSSION

Overall, the 2017 growing season was challenging for growing spring barley. The cooler than average temperatures along with the higher than normal rainfall in throughout much of the growing season created the ideal conditions for Fusarium growth. This is evident in the high DON concentrations in both varieties.

Even though all of the variety+fungicide+timing treatments resulted in DON concentrations above 1 ppm, it's important to note that Conlon, a moderately resistant variety, had lowest incidence of FHB and DON levels, while Rasmussen, a susceptible variety, had DON levels almost double (8.29 ppm) that of Conlon (4.53 ppm). This indicates the importance of selecting resistant cultivars to manage FHB in our region.

The application of the conventional fungicides Prosaro and Caramba, applied at heading, reduced DON concentrations. However, the untreated control also had low DON concentrations; this could be attributed to these plots not being sprayed with Fusarium spores (40,000 spores per ml) indicating the impact of high Fusarium inoculum during plant heading and flowering. In general, the fungicide applications at heading resulted in lower DON concentrations than the fungicides applied 4-days after heading. Interestingly, yields did not vary significantly between fungicide type, application or variety.

It is important to remember that the results only represent one year of data. The Northwest Crops and Soils Program will be repeating this trial again in 2018.

ACKNOWLEDGEMENTS

The UVM Extension Northwest Crops and Soils Team would like to thank the Borderview Research Farm for their generous help with the trials, as well as acknowledge the U.S. Wheat and Barley Scab Initiative program for their financial support. We would like to acknowledge Kelly Drollette, Hillary Emick, Abha Gupta, Freddy Morin, Lindsey Ruhl, Matt Sanders, Stuart Wolff-Goodrich and Sara Ziegler for their assistance with data collection and entry. This informatTm0 g0 G7-ET3cide applications