2019 The Efficacy of Spraying Organic Fungicides to Control Fusarium Head Blight Infection in Spring Wheat



Dr. Heather Darby, UVM Extension Agronomist
Hillary Emick and Haley Jean
UVM Extension Crops and Soils Technicians
(802) 5246501

Visit us on the web at http://www.uvm.edu/nwcrops

© December 2019, University of Vermont Extension

Glenn is a FHB resistant variety of hard red spring wheat while Shelly is moderately resistant. The fungicide+timing treatments are listed in Table 2. Listed below are the descriptions of the fungicide treatments provided by the manufacturer.

Table 2. Plot treatments-fungicide application rates.

Treatments	Application rate

Table 3. Fungicide treatment application dates, 2019.

Variety and treatment	Application date
Glenn Inoculated with *Fusarium*	24-Jun
Glenn Heading Applications	24-Jun
Glenn Post-heading Applications	29-Jun
Shelly Inoculated with *Fusarium*	29-Jun
Shelly Heading Applications	29-Jun
Shelly Post-Heading Applications	2-Jul

Grain plots were harvested with an Almaco SPC50 plot combine on 2 Aug. Following harvest, seed was cleaned with a small Clipper cleaner (A.T. Ferrell, Bluffton, IN). A one-pound subsample was collected to determine DON concentration. At the time of harvest, grain moisture and test weight were recorded with a DICKEY-John M20P meter, and yield was measured on a pound scale. Generally, the heavier the wheat is per volume, the higher baking quality. The acceptable test weight for bread wheat is approximately 56-60 lbs bu^{-1}. Subsamples were ground into flour using a Perten LM3100 Laboratory Mill in order to be evaluated for mycotoxin levels. Deoxynivalenol (DON) analysis was conducted with the Veratox DON 5/5 Quantitative test from the NEOGEN Corp., which has a detection range of 0.5 to 5 ppm. Samples with DON values greater than 1 ppm are considered unsuitable for human consumption

RESULTS

Seasonal precipitation and temperature recorded at a weather station at Borderview Research Farm are

Figure 1. The impact of application timing and fungicide on spring wheat yield and DON concentration. Treatments with the same letter did not differ significantly by DON concentration (capital letters) or yield (lowercase).

Impact of Variety

There were significant differences between varieties in test weight, with Shelly having a significantly higher test weight at 56.1 lbs bu^{-1}. Glenn had a higher yield and Shelly had a lower DON concentration, but these differences were not statistically significant.

Table 6. The impact of spring wheat variety of quality and yield, Alburgh, VT, 2019.

Variety	Harvest moisture	Test weight	Yield @13.5% moisture	DON
	%	lbs bu^{-1}	lbs ac^{-1}	ppm
Glenn	**20.8**	55.2*	**3561**	0.172
Shelly	20.9	**56.1***	3370	**0.170**
LSD (0.10)	NS	0.80	NS	NS
Variety Mean	20.9	55.7	3466	0.171

*Treatments with an asterisk are not significantly different than the top performer in bold.