2011 SPRING WHEAT VARIETY TRIAL

In 2011, the University of Vermont Extension in collaboration with the University of Maine began the second year of extensive organic variety trials evaluating hard red spring wheat in order to determine which varieties thrive in our northern climate. The trials were established at the Borderview Research Farm in Alburgh, Vermont and at Cornell University's Willsboro Research Farm in Willsboro, New York. This trial is one of several in a USDA Organic Research Education Initiative grant focused on the production of high quality organic bread wheat in New England.

MATERIALS AND METHODS

The experimental plot design was randomized complete block with four replications. Spring wheat varieties evaluated and their sources are listed in Table 1.

time flour was evaluated for its protein content, falling number, and mycotoxin levels. Grains were analyzed for protein content using the Perten Inframatic 8600 Flour Analyzer. Grain protein affects gluten strength and loaf volume. Most commercial mills target 11–15% protein. The determination of falling number (AACC Method 56-81B, AACC Intl., 2000) was

Table 4. Relative flowering date of spring wheat.

	Alburgh, VT	Willsboro, NY
Variety	Flowering date	Flowering date
07SW04		

severity in our fields. Wheat varieties differed significantly in wheat biomass at both locations (Table 5; 6). In general, the wheat dry matter yields were lower than past years; this could be attributed to the lack of wheat tillering. Interestingly, high biomass wheat varieties at Alburgh were rarely high biomass varieties at Willsboro. This may indicate that some varieties are better adapted to production in the heavy clay soils at Willsboro.

Table 5. Plant heights & weed biomass, Alburgh, VT

Alburgh,VT	

Table 6. Plant heights & weed biomass, Willsboro, NY

Willsboro, NY

Figure 1. Yields of 30 spring wheat varieties

Spring Wheat Quality:

The common measures used by commerial mills to evaluate wheat quality are: grain protein, falling number, test weigh and mycotoxin (DON) content. In Alburgh, the variety with the highest protein content was Robin, 14.7% (Table 9 and Figure 3). Other high protein varieties at this location were Helios and Kelby. In Willsboro, 07SW04 had the highest protein level at 16.6% (Table 10 and Figure 4). Robin was another high protein variety at this location. In Alburgh O7SW04 had the highest falling number, 413 seconds, and in Willsboro Ada and Helios had the highest at 473 seconds. Both Ada and Helios were amoung the varieties with the

mycotoxins. A vomitoxin called deoxynivalenol (DON) is considered the primary mycotoxin associated with FHB. The spores are usually transported by air currents and can infect plants at flowering through grain fill. Eating contaminated grain, >1ppm, poses a health risk to both humans and livestock.