



International Association for Ecology

Body-Size Differences in a Colonizing Amphipod-Mollusc Assemblage

Author(s): N. J. Gotelli, F. G. Lewis, III, C. M. Young

Source: *Oecologia*, Vol. 72, No. 1 (1987), pp. 104-108

Published by: Springer in cooperation with International Association for Ecology

Stable URL: <http://www.jstor.org/stable/4218234>

Accessed: 29/01/2010 14:17

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=springer>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Springer and International Association for Ecology are collaborating with JSTOR to digitize, preserve and extend access to *Oecologia*.




Table 3. Significant deviations from Model I, which randomized to the smallest individual (RANGE) was also smaller than

